



World Journal of Stem Cells

Submit a Manuscript: <https://www.f6publishing.com>

World J Stem Cells 2019 October 26; 11(10): 891-903

DOI: [10.4252/wjsc.v11.i10.891](https://doi.org/10.4252/wjsc.v11.i10.891)

ISSN 1948-0210 (online)

META-ANALYSIS

Stem cell treatment and cerebral palsy: Systemic review and meta-analysis

Simone Eggenberger, Céline Boucard, Andreina Schoeberlein, Raphael Guzman, Andreas Limacher, Daniel Surbek, Martin Mueller

ORCID number: Simone Eggenberger (0000-0001-6110-4952); Céline Boucard (0000-0002-6724-2392); Andreina Schoeberlein (0000-0002-6716-9551); Raphael Guzman (0000-0002-5816-3529); Andreas Limacher (0000-0002-9094-9476); Daniel Surbek (0000-0001-7705-1584); Martin Muller (0000-0001-5594-0532).

Author contributions: Eggenberger S and Mueller M drafted and wrote the manuscript; Limacher A performed the statistical analysis; Guzman R, Boucard C, Schoeberlein A, Limacher A and Surbek D revised and provided critical and important intellectual content; all authors have given final approval of the version to be published.

Conflict-of-interest statement: The authors deny any conflict of interest.

PRISMA 2009 Checklist statement: The authors have read the PRISMA 2009 Checklist, and the manuscript was prepared and revised according to the PRISMA 2009 Checklist.

Open-Access: This article is an open-access article which was selected by an in-house editor and fully peer-reviewed by external reviewers. It is distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works

Simone Eggenberger, Daniel Surbek, Martin Mueller, Department of Obstetrics and Gynaecology, Inselspital, University Hospital Bern, Bern 3010, Switzerland

Céline Boucard, Andreina Schoeberlein, Department for BioMedical Research, University of Bern, Bern 3008, Switzerland

Raphael Guzman, Department of Neurosurgery, University Hospital of Basel, Basel 4056, Switzerland

Andreas Limacher, CTU Bern, University of Bern, Bern 3012, Switzerland

Corresponding author: Martin Mueller, MD, PhD, Senior Consultant, Department of Obstetrics and Gynaecology, Inselspital, University Hospital Bern, Friedbühlstrasse 19, Bern 3010, Switzerland. martin.mueller@insel.ch

Telephone: +41-31-6321125

Fax: +41-31-6321128

Abstract

BACKGROUND

Perinatal complications may result in life-long morbidities, among which cerebral palsy (CP) is the most severe motor disability. Once developed, CP is a non-progressive disease with a prevalence of 1-2 per 1000 live births in developed countries. It demands an extensive and multidisciplinary care. Therefore, it is a challenge for our health system and a burden for patients and their families. Recently, stem cell therapy emerged as a promising treatment option and raised hope in patients and their families.

AIM

The aim is to evaluate the efficacy and safety of stem cell treatment in children with CP using a systematic review and meta-analysis

METHODS

We performed a systematic literature search on PubMed and EMBASE to find randomized controlled clinical trials (RCT) investigating the effect of stem cell transplantation in children with CP. After the review, we performed a random-effects meta-analysis focusing on the change in gross motor function, which was quantified using the gross motor function measure. We calculated the pooled standardized mean differences of the 6- and/or 12-mo-outcome by the method of Cohen. We quantified the heterogeneity using the I-squared measure.

on different terms, provided the original work is properly cited and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>

Manuscript source: Invited manuscript

Received: March 12, 2019

Peer-review started: March 15, 2019

First decision: June 3, 2019

Revised: June 12, 2019

Accepted: August 20, 2019

Article in press: August 20, 2019

Published online: October 26, 2019

P-Reviewer: Khan I, Ventura C

S-Editor: Yan JP

L-Editor: A

E-Editor: Wu YXJ



RESULTS

We identified a total of 8 RCT for a qualitative review. From the initially selected trials, 5 met the criteria and were included in the meta-analysis. Patients' population ranged from 0.5 up to 35 years ($n = 282$). We detected a significant improvement in the gross motor function with a pooled standard mean difference of 0.95 (95% confidence interval: 0.13-1.76) favoring the stem cell group and a high heterogeneity ($I^2 = 90.1\%$). Serious adverse events were rare and equally distributed among both intervention and control groups.

CONCLUSION

Stem cell therapy for CP compared with symptomatic standard care only, shows a significant positive effect on the gross motor function, although the magnitude of the improvement is limited. Short-term safety is present and further high-quality RCTs are needed.

Key words: Cerebral palsy; Perinatal brain injury; Stem cells; Umbilical cord blood; Mesenchymal stem/stromal cells; Gross motor function; Meta-analysis

©The Author(s) 2019. Published by Baishideng Publishing Group Inc. All rights reserved.

Core tip: Cerebral palsy is a severe motor disability resulting from perinatal complications. Recently, stem cell therapy emerged as a promising treatment option and raised hope in patients and their families. However, high-quality randomized clinical trials investigating the efficacy of stem cell therapy as a treatment for cerebral palsy are scarce. We included the small number of currently available trials in our meta-analysis. A slight but significantly positive effect on the gross motor function favoring the stem cell group can be seen.

Citation: Eggenberger S, Boucard C, Schoeberlein A, Guzman R, Limacher A, Surbek D, Mueller M. Stem cell treatment and cerebral palsy: Systemic review and meta-analysis. *World J Stem Cells* 2019; 11(10): 891-903

URL: <https://www.wjgnet.com/1948-0210/full/v11/i10/891.htm>

DOI: <https://dx.doi.org/10.4252/wjsc.v11.i10.891>

INTRODUCTION

Despite advances in perinatal medicine, many infants continue to face serious risks during pregnancy, parturition, and adaptation after birth. Preterm birth and severe birth asphyxia are the most frequent complications, which may lead to brain damage of the newborn. The clinical presentation in an individual child after perinatal complications is complex. This complexity results from multiple potential causal pathways, signs and symptoms of injury. Typical pathology in these newborns include brain injury and the resulting cerebral palsy (CP). Therefore, CP summarizes a heterogeneous group of non-progressive disabilities in motor function which range from slight motor impairment of an isolated body part to the inability of walking or speaking^[1]. Depending on the severity, typical symptoms are difficulties in coordination, stereotypic movement, impossibility of discrete movements, evocation of primitive reflexes through voluntary movements or co-contraction of agonist and antagonist resulting in spasticity. Multiple other impairments such as seizure disorders, altered sensation or perception or musculoskeletal disorders can appear as additional symptoms^[1]. Although the diagnosis of CP does not require cognitive disability, about two-thirds of children with CP are confronted with it^[2]. Not surprisingly, CP is the most common motor disability in children with a prevalence of 2 per 1000 live births in developed countries. The prevalence of CP increases exponentially in preterm infants with decreasing gestational age^[3]. For example, the risk to develop CP is 30 times higher in infants born before 33 wk gestation compared to term-born infants^[3]. Besides prematurity, major risk factors include placental abnormalities, major and minor birth defects, low birthweight, meconium aspiration, emergency caesarean section, birth asphyxia, neonatal seizures, respiratory distress syndrome, hypoglycemia, and neonatal infections^[4]. Together, CP originates from a multifactorial pathology with multiple risk factors and in many cases, a distinct cause

is unclear.

Due to major improvements in neonatal care, half of the children suffering from CP are preterm infants and the other half are term-born infants^[5,6]. This differentiation is important as prophylactic/therapeutic approaches differ in these two populations. Currently, the only intervention known to reduce the burden of CP in the term population is hypothermia. Hypothermia is associated with a significant reduction in death and disability in children subjected to perinatal asphyxia^[7]. However, 40%-50% of infants treated with hypothermia still die or develop significant neurological disability^[8]. In the preterm population, hypothermia is contra-productive and therapeutic options are lacking. One option is the antenatal magnesium sulfate prophylaxis at less than 30 wk of gestation, which reduces CP and combined CP and mortality rate at 2-years of age. However, long-term neurological benefits are lacking^[9,10]. New avenues to treat CP emerged and stem cell treatments are particularly promising. We review the potential of a stem cell transplant using a meta-analysis to evaluate gross motor function after randomized controlled trials (RCTs) in children with CP.

Which source of stem cells to use?

Stem cells are characterized as cells with the capacity of self-renewal and differentiation into multiple tissues^[11]. Their program of division and differentiation is regulated by the immediate microenvironment, also called the niche. Stem cells are grouped depending on the number of tissues they can differentiate into^[12]. Totipotent cells can differentiate into any cell type found in an organism. They exist for a very limited time only in the embryo shortly after fertilization. Pluripotent cells are the next stage; they occur in the blastocyst and can form cells from each of the three germinal sheets. Multipotent stem cells can also be detected in adults. Their differentiation capability is restricted to cell types within one germinal sheet. Unipotent stem cells are responsible for the renewal of a single tissue lineage. *In vitro*, apart from their self-renewal and differentiation capacity, stem cells can be identified by markers, which are gene products expressed by specific types of stem cells. Therefore, by means of flow cytometry, stem cells can be identified and quantified^[11].

Embryonic or pluripotent stem cells seem to be very promising due to their differentiation ability^[11]. However, their unlimited self-renewal and differentiation capacity combined with a lack of cell-cell-interaction and regulation through extraembryonic cells may lead to tumor formation^[13]. Additionally, the harvesting of pluripotent cells results in the death of the embryo raising major ethical concerns. Therefore, the main stem cell sources in CP treatment are bone marrow (BM) or umbilical cord blood (UCB)-derived stem cells.

Umbilical cord blood and bone marrow as a source of stem cells

UCB is a rich source of stem cells. It can be collected after birth and stored in public or private banks for a possible future use^[14]. It contains several types of stem and progenitor cells, among which hematopoietic stem cells (HSC), mesenchymal stem/stromal cells (MSC) and endothelial progenitor cells (EPC) are the most relevant. Further, the existence of embryonic-like stem cells is controversially discussed. Besides the simple isolation without ethical concerns, these cells have a remarkably low immunogenicity. Compared to stem cells from other sources, UCB cells tolerate more human leukocyte antigen (HLA)-mismatches without rejection^[12]. The reason for this might be the immature fetal immune system. Today, cord blood is routinely used to treat hematopoietic or immunologic disorders^[15]. In the last years, promising trials have shown that UCB stem cells have great potential in the treatment of various neurological diseases^[16]. A world-wide network of cord blood banks is available for unrelated cord blood transplantations. For therapeutic use, the mononuclear fraction is isolated from cord blood by means of a density gradient^[17]. The mononuclear fraction includes immunosuppressive cells such as regulatory T-cells and monocyte-derived suppressor cells^[16]. Each of these cell types has characteristics that are likely to contribute to neuroprotection. Notably, the composition of those cells depends on the timing of sampling (gestational age of pregnancy)^[18]. For example, UCB derived from preterm placentae is different in its mononuclear fraction from term-derived, and UCB from intrauterine growth restriction infants has impaired EPC^[19].

BM is the major hematopoietic organ localized in the central cavities of axial and long bones^[20]. Differentiation and proliferation of the blood cells occur in the hematopoietic compartment. It is composed of HSC and the precursors for the different blood lineages. It used to be the main source of HSC for clinical use, until recently less invasive techniques allowed to obtain HSC from other sources such as granulocyte-colony stimulating factor-stimulated peripheral blood or UCB^[21]. Importantly, the stroma is responsible for the regulation of the hematopoietic process

and contains MSC and EPC along with their products of differentiation like fibroblasts and endothelial cells. Compared to UCB, MSC are more frequent in BM. Although BM harvesting requires an invasive procedure, BM is still the main source for MSC^[22]. Notably, the number of both EPC and MSC in BM decline with age^[20,23].

Mesenchymal stem/stromal cells

MSC are a heterogeneous population of multipotent cells that can differentiate into bone (osteoblasts), fat (adipocytes), cartilage (chondroblasts) and periosteum (fibroblasts)^[24]. Their differentiation potential into the neurogenic direction is debated, but the discussion remains controversial^[17]. Originally, MSCs were discovered in BM as part of the mononuclear cell fraction, where they support HSC^[25]. Meanwhile, they have been isolated from many other sources including adipose tissue, muscle, placental tissues, and UCB. The International Society for Cell and Gene Therapy elaborated minimal criteria to define MSCs^[26]. It remains unclear and much discussed what MSC's function in each source is. Some even consider the pericyte to be the cell of origin of MSC *in vivo*. Thus, their function could be the regulation of the capillary blood flow and permeability^[27]. There are high expectations for MSC as a therapy in various diseases. A special interest lies in the treatment of neurological diseases including CP^[24]. Also, MSC have become more and more commercialized as source of replacement for damaged structures. However, injected allogeneic MSC are rejected by the host immune system and likely to be eliminated soon after the transplantation and in contrast, autologous ones may persist for some longer time^[27]. Not surprisingly, the therapeutic effect of MSC is attributed not to the differentiation capacity and thus formation of new tissue, but to the secretome, which contains modulatory factors^[28]. These modulate oxidative stress and has angiogenetic, anti-apoptotic and anti-inflammatory effects^[28]. MSC have different characteristics depending on their origin. Of special interest are those MSC derived from fetal tissues such as UCB, placental or cord tissues, which are believed to have a wider differentiation and greater proliferation potential^[29,30]. Moreover, they can easily be isolated non-invasively, rapidly and without ethical concern nor invasive procedures^[31]. However, we are not always successful to isolate a sufficient number of MSC from UCB for clinical use^[32].

Hematopoietic stem cells and endothelial progenitor cells

Both HSC and EPC can be found in BM as well as in UCB. HSC are multipotent precursor cells with the ability to form all blood cells while maintaining a sufficient HSC-pool to provide hematopoiesis throughout life^[33]. Whilst HSC transplantation was the first established stem cell transplantation and exclusively for hematologic disorders, HSC now also become interesting in the field of non-hematologic diseases^[34]. Of special interest for a possible neuroprotective contribution is their beneficial effect in ischemic brain injury in animal models^[35]. EPC too are believed to possess neuroprotective features. Apart from their ability to differentiate into endothelial cells, they are believed to induce neovascularization and reduce hypoxia-induced apoptosis and destruction of blood vessels^[36,37]. Considering these in-vitro characteristics of both HSC and EPC, they may play a role in neuroprotective therapies such as the treatment of CP as well.

MATERIALS AND METHODS

Literature search

We followed the PRISMA-statement guidelines to perform a systematic electronic search using PubMed and EMBASE databases for trials published between January 1, 1990, and February 11, 2019. To be eligible for inclusion, the trial was required to be a randomized, controlled clinical trial with full-text availability in English. Further, the study population must consist of children diagnosed with any type of CP. The intervention must include any kind of stem cell treatment compared to placebo and/or standard of care such as rehabilitation. The motor outcome of both intervention and control group must be assessed and reported in the Gross Motor Function Measure (GMFM). We excluded trials that did not meet these criteria from the analysis.

We combined terms like “stem cells”, “neural progenitor cells”, “mesenchymal stem cells” or “umbilical cord blood” to match the intervention with “cerebral palsy” in both databases. We set the limitations to human-controlled clinical trials in English only. Also, we detected trials from bibliographies of other articles. We identified the potentially relevant articles through title and abstract screening and proceeded to the definite inclusion or exclusion according to their content (supplementary materials). The study selection process is illustrated in [Figure 1](#).




Figure 1 Study selection process.

Meta-analysis

We defined the efficacy of stem cell therapy as a change in gross motor function. To summarize the gross motor outcome from the intervention group compared to the control group, we performed a random-effects meta-analysis using the method of DerSimonian *et al.*^[38]. We pooled the standardized mean differences and the heterogeneity was quantified using the I-squared measure, taken from the inverse-variance fixed-effect model. All analyses were done in Stata version 15 using the command meta version 3.04.

RESULTS

Study selection and characteristics

After applying the inclusion and exclusion criteria, we identified 8 relevant trials for further analysis (Figure 1). Four of them have been conducted in China, three in Korea and one in the United States of America. Publication dates ranged from 2012 to 2018. The study characteristics are summarized in Table 1. Five trials investigated stem cell therapy plus rehabilitation to rehabilitation only, one trial compared stem cell therapy only to rehabilitation and two trials compared stem cell therapy only to no intervention in the control group. Two of the included studies were three-group randomized clinical trials investigating the additive effect of erythropoietin^[39] or using mononuclear cells^[40]. Two other studies were designed as randomized controlled crossover-trials^[41,42]. The age in the study population ranged from 6 mo to 35 years. All patients included were diagnosed with CP. The severity of CP was measured with the Gross Motor Function Classification System, which divides the syndrome of CP into 5 levels, whereof level 5 is the most severe motor impairment^[43]. The type and dose of the stem cells differed in the trials. One trial used neural progenitor cells derived from aborted fetuses' forebrains^[44], while others included umbilical cord- or BM-derived stem cells^[39,41,45-47]. The applied dose ranged from 4×10^6 to 6×10^8 cells. The number of cells was not always adapted to body weight. The application route was in most cases the intravenous route. In summary, both the study population and the intervention characteristics are heterogeneous, which leads to a certain risk of bias summarized in Table 2 and Figure 2.

Risk of bias

We assessed the risk of bias in individual studies using the Cochrane criteria. The risk ranged from low to high. The major source of risk of bias was patient and personnel blinding^[40,44-46]. The number of trials was not sufficient to estimate the risk for publication bias.

Outcome measurements

All included trials applied standardized scales to assess the neurodevelopmental

Table 1 Intervention and study population overview

Ref.	Country	Study Population		Patient condition	Intervention					
		Sample size	Age (yr)		Intervention on and study groups	Application route	Type of stem cells	Source	Origin	Dose
Chen <i>et al</i> ^[45] , 2013	China	60	1-35	CP, GMFCS level 3 to 5	Group 1: SC + rehabilitation Group 2: Rehabilitation only	Intrathecal	Mesenchymal stem cells <i>in vitro</i> transformed to neural stem cell-like cells	Bone marrow	Autologous	Fixed quantity of $1-2 \times 10^7$ cells, twice
Huang <i>et al</i> ^[46] , 2018	China	56	3-12	CP	Group 1: SC + rehabilitation Group 2: Placebo + rehabilitation	Intravenous	Mesenchymal stem cells	Cord blood	Allogeneic	Fixed quantity of 5×10^7 cells, 4 times
Kang <i>et al</i> ^[47] , 2015	South Korea	36	0.5-20	CP, GMFCS level 1 to 5	Group 1: SC Group 2: Placebo No rehabilitation	Intravenous/intra-arterial	Mononuclear cells	Cord blood	Allogeneic ¹	5.46×10^7 cells/kg, once
Liu <i>et al</i> ^[40] , 2017	China	105	0.5-12.5	Spastic CP, GMFCS level 2 to 5	Group 1: MNC Group 2: MSC Group 3: rehabilitation	Intrathecal	Mesenchymal stromal cells and mononuclear cells	Bone marrow	Autologous	10^6 cells/kg, 4 times
Luan <i>et al</i> ^[44] , 2012	China	91	< 0.5-3.5	CP, GMFCS level 1 to 5	Group 1: SC + rehabilitation Group 2: rehabilitation only	Intraventricular	Neural progenitor cells	Aborted healthy fetuses' forebrains	Allogeneic	Fixed quantity of $8-10 \times 10^6$ cells, once
Min <i>et al</i> ^[39] , 2013	South Korea	105	0.8-10	CP	Group 1: SC+ EPO + rehabilitation Group 2: placebo SC + EPO + rehabilitation Group 3: placebo SC+ placebo + rehabilitation	Intravenous	Mononuclear cells	Cord blood	Allogeneic ¹	$\geq 3 \times 10^7$ cells/kg, once
Rah <i>et al</i> ^[42] , 2017	South Korea	57	2-10	Non-severe CP	Group 1: G-CSF at baseline, SC 1 mo later Group 2: G-CSF at baseline, placebo 1 mo later crossover at 7 mo	Intravenous	Mobilized mononuclear cells	Peripheral blood	Autologous	5.97×10^8 cells/kg, once

Sun <i>et al.</i> ^[41] , 2017	USA	63	1-6	CP of any type, GMFCS Level 2 to 4 or 1 with hemiplegia	Group 1: SC + rehabilitation Group 2: Placebo + rehabilitation	Intravenous	Mononuclear cells	Cord blood	Autologous	2 × 10 ⁷ cells/kg, once
--	-----	----	-----	---	--	-------------	-------------------	------------	------------	------------------------------------

¹Administration of cyclosporine. MNC: Mononuclear cells; MSC: Mesenchymal stem cells; EPO: Erythropoietin; CP: Cerebral palsy; SC: Stem cells; GMFCS: Gross motor function classification system.

outcome. The fine and gross motor function, cognitive function and spasticity were measured. In the following analysis, we focused on the gross motor function outcome only. The gross motor function was evaluated with the GMFM-66 or GMFM-88^[43]. Both scores categorize 5 sections: Lying, Sitting, Crawling, Standing and Walking. Each of these categories is expressed in the percentage of the maximum achievable points. To determine the total score, the percentages of all categories are added up and divided by five and therefore expresses the average score of all categories. Importantly, the total score varies depending on the severity of CP and the patient's age. The observation period in the included trials was 6 months in three trials, 12 mo in another three and 24 in the remaining two trials (see Table 3 for details). Functional assessments to measure the outcomes were performed at baseline, after the observation period, and additionally at one to three points in-between. Most GMFM data were available for the 6- and 12-mo periods and therefore we decided to proceed with those for the meta-analysis.

Cell dose, gross motor function and safety

Three trials from the initial literature search had to be excluded from the meta-analysis due to the lack of GMFM scores in an averaged format^[45], missing standard deviation or standard error in the GMFM scale^[42], or no comparable outcome score of the control group given^[44]. One study did not provide changes in GMFM summary scores from baseline, but only absolute GMFM dimension scores at baseline, 6 months and 12 mo^[40]. Therefore, we averaged the dimension scores and their variations and calculated the changes from the baseline. The cell dose used in the included trials differed but improved motor outcome seems to correlate with a higher cell dose^[41,42,47] (Tables 1 and 3). We generated a forest plot with the available GMFM-scores at 6 and 12 mo (see Figures 3-5). Additionally, we pooled all studies; the resulting forest plot is presented in Figure 5. Besides the improvement of gross motor function, the studies assessed the serious adverse events (SAE). Overall, only two trials reported the occurrence of SAEs^[39,44] (Table 4). One case of hemorrhagic foci after transplantation was detected and was presumably linked to the invasive procedure^[44]. The other SAEs mainly consisted of infections or seizures^[39]. These SAEs were equally distributed between both cell recipient and control groups. There is no report of an overdose in any of the included trials. The highest dose applied was a mean dose of 5.97×10^8 cells/kg and no SAEs were reported^[42]. Together, the included RCTs provide short-term safety, whereas the long-term impact is unclear.

DISCUSSION

Using a systematic literature review, we identified 8 RCTs investigating the effect of stem cell treatment in children with CP *vs* standard care with or without placebo. Based on our inclusion criteria, we included 5 trials in the meta-analysis with a total of 282 patients (142 control, 140 cell recipient group). By combining the outcome, we identified a significant improvement in the cell recipient compared to the control group (Figure 5). The pooled standard mean difference was 0.95 (95% confidence interval: 0.13-1.76). However, we detected a high heterogeneity ($I^2 = 90.1\%$), which reflects the diverse patient characteristics and needs to be accounted for while discussing the results. Besides, our analysis only includes the results from 5 trials, which is hardly enough to represent the whole CP population. A search on clinicaltrials.gov identified 9 ongoing randomized clinical trials about stem cell therapy in CP. Therefore, evidence might grow stronger in the future with greater data acquisition.

In the included trials, the origin (autologous or allogeneic), type (MSC, mononuclear cells, and neural progenitor cells), and source (BM, brain tissue, UCB) of the transplanted cells differed. Interestingly, in patients receiving the transplant with

Table 2 Evaluation of the risk of bias

Ref.	Random sequence generation: Selection bias	Allocation concealment	Blinding of participants and personnel: Performance bias	Blinding of outcome assessment: Detection bias	Incomplete outcome data: Attrition bias	Selective reporting
Chen <i>et al</i> ^[45] , 2013	High	High	High	Low	High	High
Huang <i>et al</i> ^[46] , 2018	Unclear	Unclear	High	Low	Low	Unclear
Kang <i>et al</i> ^[47] , 2015	Low	Low	Low	Low	Low	Unclear
Liu <i>et al</i> ^[40] , 2017	Low	Unclear	High	Unclear	Low	Low
Luan <i>et al</i> ^[44] , 2012	Unclear	Unclear	High	Unclear	High	Unclear
Min <i>et al</i> ^[39] , 2013	Low	Low	Low	Low	Low	Low
Rah <i>et al</i> ^[42] , 2017	Low	Low	Low	Low	High	Unclear
Sun <i>et al</i> ^[41] , 2017	Low	Low	Low	Low	Low	Unclear

fewer HLA-mismatches, the improvement of gross motor function was clearly evident^[39,47]. We speculate that autologous cells may show a greater benefit in motor function improvement. Furthermore, the matching of HLA seems to have less relevance in case of a MSC transplant^[46]. For example, both studies with a significant increase in the motor outcome used a MSC transplant^[40,46]. These observations are further supported by the direct comparison of mononuclear fraction of BM stem cells and MSC alone^[40], which clearly favors a MSC alone approach. Together, the present studies suggest that MSC are the ideal candidates to modulate the gross motor function in CP. The source may be the UCB as it is less invasive and easily accessible. However, harvesting a sufficient number of cells can only be achieved in around 40% of the samples^[48], which limits the use of autologous UCB-derived MSC. Further studies are warranted especially as other easily available MSC sources such as Wharton's jelly of the umbilical cord are very promising^[49,50].

In order to report a proper gross motor function improvement, both the patient's age and level of motor impairment need to be considered by using percentiles^[51]. However, only one of the included trials used proper percentiles and presented the results as "greater than expected"^[41]. Interestingly, this trial detected a significant gross motor function improvement using a higher cell dose ($\geq 2 \times 10^7$ cells) only. Another consideration is the use of repeated cell applications^[40,46]. Given that the pathophysiology of CP consists of persistent inflammation, such repetitive doses would be plausible^[52]. Moreover, the transplanted cells do not persist for a very long time^[53], which further supports concurrent use. Eventually, the included trials do not clearly provide a specific effective dose of cells, but it should be more than 2×10^7 total nucleated cells and may go up to 6×10^8 without the risk of overdosing^[41,42]. The invasive transplantation routes such as the lumbar puncture were chosen to bypass the blood-brain barrier but resulted in a higher number of reactions like nausea and headache^[40]. However, the modulation of inflammatory responses by secretion of factors can be achieved peripherally^[54], making the route of application less relevant for the efficacy of the cells^[55].

In conclusion, stem cell therapy for CP compared with symptomatic standard care only, shows a significant positive effect on the gross motor function, although the magnitude of the improvement is limited. Especially MSC show a positive outcome. This novel treatment seems safe, at least in the short term. However, high-quality RCTs are still lacking.


Table 3 Raw motor outcome data

Ref.	Follow-up assessments after intervention (mo)	Change in GMFM-88/66 after 6 mo \pm SD	Change in GMFM-88/66 after 12 mo \pm SD	% improvement compared to baseline after whole observation period
Chen <i>et al</i> ^[45] , 2013	1, 3, 6	Numbers inaccurately indicated	-	-
Huang <i>et al</i> ^[46] , 2018	3, 6, 12, 24	Control: + 2.96 \pm 1.66; MSC: + 7.62 \pm 2.44	Control: + 4.75 \pm 1.45; MSC: + 10.27 \pm 2.96	Control: 5.67%; UCB: 14.89%
Kang <i>et al</i> ^[47] , 2015	1, 3, 6	Control: 3.85 \pm 3.75; UCB: 7.08 \pm 7.36		Control: 13.1%; UCB: 24.2%
Liu <i>et al</i> ^[40] , 2017	3, 6, 12	Control: + 1.85; BMMNC: + 3.1; BMMSC: + 10.45	Control: + 2.91; BMMNC: + 6.46; BMMSC: + 12.52	Control: 7.84%; BMMSC: 33.76%
Luan <i>et al</i> ^[44] , 2012	1, 6, 12	Control: not indicated; NPC: + 8.86	-	-
Min <i>et al</i> ^[39] , 2013	1, 3, 6	Control: + 7.8 \pm 5.1; EPO only: + 9.0 \pm 6.3; UCB + EPO: + 9.1 \pm 6.7	-	No baseline score provided
Rah <i>et al</i> ^[42] , 2017	12 mo after treatment resp. 6 mo for crossover-group	-	Transplantation at baseline: + 2.9; Transplantation after 6 mo: + 6.37	No baseline score provided
Sun <i>et al</i> ^[41] , 2017	12, 24	-	Control: + 6.9 \pm 5.5; UCB: + 7.5 \pm 6.8 (High dose: improvement + 4.3 \pm 1.5 greater than expected. Low-dose and placebo: no significant improvement beyond expectation.)	Control: 13.3%; UCB: 15.3%

BMMNC: Bone marrow mononuclear cells; BMMSC: Bone marrow mesenchymal stem cells; EPO: Erythropoietin; NPC: Neural progenitor cells; UCB: Umbilical cord blood; MSC: Mesenchymal stem cells; GMFM: Gross Motor Function Measure.

Table 4 Reported serious adverse events

Ref.	Serious adverse events in cell recipient group	Serious adverse events in control group	Allogenic cells	Application route
Chen <i>et al</i> ^[45] , 2013	0/30	0/30		Intrathecal
Huang <i>et al</i> ^[46] , 2018	0/27	0/27	x	Intravenous
Kang <i>et al</i> ^[47] , 2015	0/18	0/18	x (+ immunosuppressant)	Intravenous/intraarterial
Liu <i>et al</i> ^[40] , 2017	0/68	0/34		Intrathecal
Luan <i>et al</i> ^[44] , 2012	1/45	0/49	x	Intraventricular
Min <i>et al</i> ^[39] , 2013	3/36	6/70	x (+ immunosuppressant)	Intravenous
Rah <i>et al</i> ^[42] , 2017	0/57	-		Intravenous
Sun <i>et al</i> ^[41] , 2017	0/63	-		Intravenous
Total	4/344 (1.2%)	6/228 (2.6%)		


Figure 2 Overall risk of bias.




Figure 3 Forest plot showing the effect size of the change in the Gross Motor Function Measure in the intervention group compared to the control group after 6 mo. GMFM: Gross Motor Function Measure; SMD: Standard mean difference; CI: Confidence interval.




Figure 4 Forest plot showing the effect size of the change in the Gross Motor Function Measure in the intervention group compared to the control group after 12 mo. GMFM: Gross Motor Function Measure; SMD: Standard mean difference; CI: Confidence interval.




Figure 5 Forest plot showing the effect size of the change in the Gross Motor Function Measure in the intervention group compared to the control group. If available, measures at 12 mo were used^[40,41,46], else, measures at 6 mo were used^[39,47]. GMFM: Gross Motor Function Measure; SMD: Standard mean difference; CI: Confidence interval.

ARTICLE HIGHLIGHTS

Research background

Cerebral palsy (CP) is a severe, incurable motor disability resulting from perinatal complications. It is a challenge for our health system and a burden for both patients and their families.

Research motivation

During the last years, stem cell therapy emerged as a novel treatment option. Clinical trials report promising results and create high expectations. However, most trials are of poor quality and neither randomized nor controlled, so the scientific evidence remains doubtful.

Research objectives

The aim of our meta-analysis was to investigate the effect of stem cell treatment on the gross motor function in children with CP.

Research methods

With a systematic literature search on PubMed and EMBASE, we identified the eligible randomized controlled clinical trials (RCTs). We performed a random-effects meta-analysis focusing on the change in gross motor function and calculated the pooled standardized mean differences of the 6- and/or 12-mo-outcome.

Research results

We identified a total of 8 RCTs for a qualitative review. From the initially selected trials, 5 met the criteria and were included in the meta-analysis. Patients' population ranged from 0.5 up to 35 years ($n = 282$). We detected a significant improvement in the gross motor function with a pooled standard mean difference of 0.95 (95% confidence interval: 0.13-1.76) favoring the stem cell group and a high heterogeneity ($I^2 = 90.1\%$). Serious adverse events were rare and equally distributed among both intervention and control groups.

Research conclusions

Stem cell therapy for CP compared with symptomatic standard care only, shows a significant positive effect on the gross motor function, although the magnitude of the improvement is limited.

Research perspectives

Considering that this small number may not be enough to represent the whole CP population, our meta-analysis detected a small but significant improvement in the gross motor function favoring the stem cell group. However, the magnitude of the effect is limited. In the future, high-quality research with a more homogenous study population is needed to bring more clarity.

REFERENCES

- 1 **Rosenbaum P**, Paneth N, Leviton A, Goldstein M, Bax M, Damiano D, Dan B, Jacobsson B. A report: The definition and classification of cerebral palsy April 2006. *Dev Med Child Neurol Suppl* 2007; **109**: 8-14 [PMID: [17370477](#) DOI: [10.1111/j.1469-8749.2007.tb12610.x](#)]
- 2 **Novak I**, Hines M, Goldsmith S, Barclay R. Clinical prognostic messages from a systematic review on cerebral palsy. *Pediatrics* 2012; **130**: e1285-e1312 [PMID: [23045562](#) DOI: [10.1542/peds.2012-0924](#)]
- 3 **Oskoui M**, Coutinho F, Dykeman J, Jetté N, Pringsheim T. An update on the prevalence of cerebral palsy: A systematic review and meta-analysis. *Dev Med Child Neurol* 2013; **55**: 509-519 [PMID: [23346889](#) DOI: [10.1111/dmcn.12080](#)]
- 4 **McIntyre S**, Taitz D, Keogh J, Goldsmith S, Badawi N, Blair E. A systematic review of risk factors for cerebral palsy in children born at term in developed countries. *Dev Med Child Neurol* 2013; **55**: 499-508 [PMID: [23181910](#) DOI: [10.1111/dmcn.12017](#)]
- 5 **Colver A**, Fairhurst C, Pharoah PO. Cerebral palsy. *Lancet* 2014; **383**: 1240-1249 [PMID: [24268104](#) DOI: [10.1016/S0140-6736\(13\)61835-8](#)]
- 6 **Van Naarden Braun K**, Doernberg N, Schieve L, Christensen D, Goodman A, Yeargin-Allsopp M. Birth Prevalence of Cerebral Palsy: A Population-Based Study. *Pediatrics* 2016; **137** [PMID: [26659459](#) DOI: [10.1542/peds.2015-2872](#)]
- 7 **Jacobs SE**, Berg M, Hunt R, Tarnow-Mordi WO, Inder TE, Davis PG. Cooling for newborns with hypoxic ischaemic encephalopathy. *Cochrane Database Syst Rev* 2013; CD003311 [PMID: [23440789](#) DOI: [10.1002/14651858.CD003311.pub3](#)]
- 8 **Edwards AD**, Brocklehurst P, Gunn AJ, Halliday H, Juszczak E, Levene M, Strohm B, Thoresen M, Whitelaw A, Azzopardi D. Neurological outcomes at 18 months of age after moderate hypothermia for perinatal hypoxic ischaemic encephalopathy: Synthesis and meta-analysis of trial data. *BMJ* 2010; **340**: c363 [PMID: [20144981](#) DOI: [10.1136/bmj.c363](#)]
- 9 **Cahill AG**, Stout MJ, Caughey AB. Intrapartum magnesium for prevention of cerebral palsy: Continuing controversy? *Curr Opin Obstet Gynecol* 2010; **22**: 122-127 [PMID: [20179596](#) DOI: [10.1097/GCO.0b013e3283372312](#)]
- 10 **Doyle LW**, Anderson PJ, Haslam R, Lee KJ, Crowther C; Australasian Collaborative Trial of Magnesium Sulphate (ACTOMgSO4) Study Group. School-age outcomes of very preterm infants after antenatal treatment with magnesium sulfate vs placebo. *JAMA* 2014; **312**: 1105-1113 [PMID: [25226476](#) DOI: [10.1001/jama.2014.11189](#)]
- 11 **Slack JMW**. What is a stem cell? *Wiley Interdiscip Rev Dev Biol* 2018; e323 [PMID: [29762894](#) DOI: [10.1002/wdev.323](#)]
- 12 **Forraz N**, McGuckin CP. The umbilical cord: A rich and ethical stem cell source to advance regenerative medicine. *Cell Prolif* 2011; **44** Suppl 1: 60-69 [PMID: [21481046](#) DOI: [10.1111/j.1365-2184.2010.00729.x](#)]
- 13 **Ben-David U**, Benvenisty N. The tumorigenicity of human embryonic and induced pluripotent stem cells. *Nat Rev Cancer* 2011; **11**: 268-277 [PMID: [21390058](#) DOI: [10.1038/nrc3034](#)]
- 14 **Carroll JE**, Mays RW. Update on stem cell therapy for cerebral palsy. *Expert Opin Biol Ther* 2011; **11**: 463-471 [PMID: [21299445](#) DOI: [10.1517/14712598.2011.557060](#)]
- 15 **Gluckman E**. History of cord blood transplantation. *Bone Marrow Transplant* 2009; **44**: 621-626 [PMID: [19344444](#) DOI: [10.1038/bmt.2009.10](#)]

- 19802032 DOI: [10.1038/bmt.2009.280](https://doi.org/10.1038/bmt.2009.280)]
- 16 **McDonald CA**, Fahey MC, Jenkin G, Miller SL. Umbilical cord blood cells for treatment of cerebral palsy; timing and treatment options. *Pediatr Res* 2018; **83**: 333-344 [PMID: [28937975](https://pubmed.ncbi.nlm.nih.gov/28937975/) DOI: [10.1038/pr.2017.236](https://doi.org/10.1038/pr.2017.236)]
- 17 **Galieva LR**, Mukhamedshina YO, Arkhipova SS, Rizvanov AA. Human Umbilical Cord Blood Cell Transplantation in Neuroregenerative Strategies. *Front Pharmacol* 2017; **8**: 628 [PMID: [28951720](https://pubmed.ncbi.nlm.nih.gov/28951720/) DOI: [10.3389/fphar.2017.00628](https://doi.org/10.3389/fphar.2017.00628)]
- 18 **Messerli M**, Wagner A, Sager R, Mueller M, Baumann M, Surbek DV, Schoeberlein A. Stem cells from umbilical cord Wharton's jelly from preterm birth have neuroglial differentiation potential. *Reprod Sci* 2013; **20**: 1455-1464 [PMID: [23670950](https://pubmed.ncbi.nlm.nih.gov/23670950/) DOI: [10.1177/1933719113488443](https://doi.org/10.1177/1933719113488443)]
- 19 **Sipos PI**, Bourque SL, Hubel CA, Baker PN, Sibley CP, Davidge ST, Crocker IP. Endothelial colony-forming cells derived from pregnancies complicated by intrauterine growth restriction are fewer and have reduced vasculogenic capacity. *J Clin Endocrinol Metab* 2013; **98**: 4953-4960 [PMID: [24106289](https://pubmed.ncbi.nlm.nih.gov/24106289/) DOI: [10.1210/jc.2013-2580](https://doi.org/10.1210/jc.2013-2580)]
- 20 **Rurali E**, Bassetti B, Perrucci GL, Zanobini M, Malafronte C, Achilli F, Gambini E. BM ageing: Implication for cell therapy with EPCs. *Mech Ageing Dev* 2016; **159**: 4-13 [PMID: [27045606](https://pubmed.ncbi.nlm.nih.gov/27045606/) DOI: [10.1016/j.mad.2016.04.002](https://doi.org/10.1016/j.mad.2016.04.002)]
- 21 **Deotare U**, Al-Dawsari G, Couban S, Lipton JH. G-CSF-primed bone marrow as a source of stem cells for allografting: Revisiting the concept. *Bone Marrow Transplant* 2015; **50**: 1150-1156 [PMID: [25915812](https://pubmed.ncbi.nlm.nih.gov/25915812/) DOI: [10.1038/bmt.2015.80](https://doi.org/10.1038/bmt.2015.80)]
- 22 **Borges FT**, Convento MB, Schor N. Bone marrow-derived mesenchymal stromal cell: What next? *Stem Cells Cloning* 2018; **11**: 77-83 [PMID: [30510433](https://pubmed.ncbi.nlm.nih.gov/30510433/) DOI: [10.2147/SCCAA.S147804](https://doi.org/10.2147/SCCAA.S147804)]
- 23 **Stolzinger A**, Jones E, McGonagle D, Scutt A. Age-related changes in human bone marrow-derived mesenchymal stem cells: Consequences for cell therapies. *Mech Ageing Dev* 2008; **129**: 163-173 [PMID: [18241911](https://pubmed.ncbi.nlm.nih.gov/18241911/) DOI: [10.1016/j.mad.2007.12.002](https://doi.org/10.1016/j.mad.2007.12.002)]
- 24 **Mukai T**, Tojo A, Nagamura-Inoue T. Mesenchymal stromal cells as a potential therapeutic for neurological disorders. *Regen Ther* 2018; **9**: 32-37 [PMID: [30525073](https://pubmed.ncbi.nlm.nih.gov/30525073/) DOI: [10.1016/j.reth.2018.08.001](https://doi.org/10.1016/j.reth.2018.08.001)]
- 25 **Hematti P**, Viswanathan S, Hematti P. Mesenchymal Stromal Cells: Clinical Experience, Challenges, and Future Directions. Viswanathan S, Hematti P. *Mesenchymal Stromal Cells*. Boston: Academic Press 2017; 309-334 [DOI: [10.1016/B978-0-12-802826-1.00012-X](https://doi.org/10.1016/B978-0-12-802826-1.00012-X)]
- 26 **Dominici M**, Le Blanc K, Mueller I, Slaper-Cortenbach I, Mueller I, Marini F, Krause D, Deans R, Keating A, Prockop DJ, Horwitz E. Minimal criteria for defining multipotent mesenchymal stromal cells. The International Society for Cellular Therapy position statement. *Cytotherapy* 2006; **8**: 315-317 [PMID: [16923606](https://pubmed.ncbi.nlm.nih.gov/16923606/) DOI: [10.1080/14653240600855905](https://doi.org/10.1080/14653240600855905)]
- 27 **Slack JM**. *Regeneration, Wound Healing and Cancer*. In: The Science of Stem cells. John Wiley and Sons, Ltd 2017; 217-238 [DOI: [10.1002/9781119235293.ch11](https://doi.org/10.1002/9781119235293.ch11)]
- 28 **Atkinson K**. *The mesenchymal stem cell, the mesenchymal stromal cell, and the mesenchymal stromal cell exosome*. In: The Biology and Therapeutic Application of Mesenchymal Cells. John Wiley and Sons, Ltd 2016; 1-7 [DOI: [10.1002/9781118907474.ch1](https://doi.org/10.1002/9781118907474.ch1)]
- 29 **van de Ven C**, Collins D, Bradley MB, Morris E, Cairo MS. The potential of umbilical cord blood multipotent stem cells for nonhematopoietic tissue and cell regeneration. *Exp Hematol* 2007; **35**: 1753-1765 [PMID: [17949892](https://pubmed.ncbi.nlm.nih.gov/17949892/) DOI: [10.1016/j.exphem.2007.08.017](https://doi.org/10.1016/j.exphem.2007.08.017)]
- 30 **Joerger-Messerli M**, Brühlmann E, Bessire A, Wagner A, Mueller M, Surbek DV, Schoeberlein A. Preeclampsia enhances neuroglial marker expression in umbilical cord Wharton's jelly-derived mesenchymal stem cells. *J Matern Fetal Neonatal Med* 2015; **28**: 464-469 [PMID: [24803009](https://pubmed.ncbi.nlm.nih.gov/24803009/) DOI: [10.3109/14767058.2014.921671](https://doi.org/10.3109/14767058.2014.921671)]
- 31 **Mukai T**, Tojo A, Nagamura-Inoue T. Umbilical Cord-Derived Mesenchymal Stromal Cells Contribute to Neuroprotection in Neonatal Cortical Neurons Damaged by Oxygen-Glucose Deprivation. *Front Neurol* 2018; **9**: 466 [PMID: [29963009](https://pubmed.ncbi.nlm.nih.gov/29963009/) DOI: [10.3389/fneur.2018.00466](https://doi.org/10.3389/fneur.2018.00466)]
- 32 **Zeddou M**, Briquet A, Relic B, Josse C, Malaise MG, Gothot A, Lechanteur C, Beguin Y. The umbilical cord matrix is a better source of mesenchymal stem cells (MSC) than the umbilical cord blood. *Cell Biol Int* 2010; **34**: 693-701 [PMID: [20187873](https://pubmed.ncbi.nlm.nih.gov/20187873/) DOI: [10.1042/CBI20090414](https://doi.org/10.1042/CBI20090414)]
- 33 **Hawley RG**, Ramezani A, Hawley TS. Hematopoietic stem cells. *Methods Enzymol* 2006; **419**: 149-179 [PMID: [17141055](https://pubmed.ncbi.nlm.nih.gov/17141055/) DOI: [10.1016/S0076-6879\(06\)19007-2](https://doi.org/10.1016/S0076-6879(06)19007-2)]
- 34 **Porada CD**, Atala AJ, Almeida-Porada G. The hematopoietic system in the context of regenerative medicine. *Methods* 2016; **99**: 44-61 [PMID: [26319943](https://pubmed.ncbi.nlm.nih.gov/26319943/) DOI: [10.1016/j.ymeth.2015.08.015](https://doi.org/10.1016/j.ymeth.2015.08.015)]
- 35 **Felfly H**, Haddad GG. Hematopoietic stem cells: Potential new applications for translational medicine. *J Stem Cells* 2014; **9**: 163-197 [PMID: [25157450](https://pubmed.ncbi.nlm.nih.gov/25157450/)]
- 36 **Tanaka N**, Kamei N, Nakamae T, Yamamoto R, Ishikawa M, Fujiwara H, Miyoshi H, Asahara T, Ochi M, Kudo Y. CD133+ cells from human umbilical cord blood reduce cortical damage and promote axonal growth in neonatal rat organ co-cultures exposed to hypoxia. *Int J Dev Neurosci* 2010; **28**: 581-587 [PMID: [20673797](https://pubmed.ncbi.nlm.nih.gov/20673797/) DOI: [10.1016/j.ijdevneu.2010.07.232](https://doi.org/10.1016/j.ijdevneu.2010.07.232)]
- 37 **Murohara T**. Therapeutic vasculogenesis using human cord blood-derived endothelial progenitors. *Trends Cardiovasc Med* 2001; **11**: 303-307 [PMID: [11728877](https://pubmed.ncbi.nlm.nih.gov/11728877/) DOI: [10.1016/s1050-1738\(01\)00128-1](https://doi.org/10.1016/s1050-1738(01)00128-1)]
- 38 **DerSimonian R**, Laird N. Meta-analysis in clinical trials. *Control Clin Trials* 1986; **7**: 177-188 [PMID: [3802833](https://pubmed.ncbi.nlm.nih.gov/3802833/) DOI: [10.1016/0197-2456\(86\)90046-2](https://doi.org/10.1016/0197-2456(86)90046-2)]
- 39 **Min K**, Song J, Kang JY, Ko J, Ryu JS, Kang MS, Jang SJ, Kim SH, Oh D, Kim MK, Kim SS, Kim M. Umbilical cord blood therapy potentiated with erythropoietin for children with cerebral palsy: A double-blind, randomized, placebo-controlled trial. *Stem Cells* 2013; **31**: 581-591 [PMID: [23281216](https://pubmed.ncbi.nlm.nih.gov/23281216/) DOI: [10.1002/stem.1304](https://doi.org/10.1002/stem.1304)]
- 40 **Liu X**, Fu X, Dai G, Wang X, Zhang Z, Cheng H, Zheng P, An Y. Comparative analysis of curative effect of bone marrow mesenchymal stem cell and bone marrow mononuclear cell transplantation for spastic cerebral palsy. *J Transl Med* 2017; **15**: 48 [PMID: [28235424](https://pubmed.ncbi.nlm.nih.gov/28235424/) DOI: [10.1186/s12967-017-1149-0](https://doi.org/10.1186/s12967-017-1149-0)]
- 41 **Sun JM**, Song AW, Case LE, Mikati MA, Gustafson KE, Simmons R, Goldstein R, Petry J, McLaughlin C, Waters-Pick B, Chen LW, Wease S, Blackwell B, Worley G, Troy J, Kurtzberg J. Effect of Autologous Cord Blood Infusion on Motor Function and Brain Connectivity in Young Children with Cerebral Palsy: A Randomized, Placebo-Controlled Trial. *Stem Cells Transl Med* 2017; **6**: 2071-2078 [PMID: [29080265](https://pubmed.ncbi.nlm.nih.gov/29080265/) DOI: [10.1002/sctm.17-0102](https://doi.org/10.1002/sctm.17-0102)]
- 42 **Rah WJ**, Lee YH, Moon JH, Jun HJ, Kang HR, Koh H, Eom HJ, Lee JY, Lee YJ, Kim JY, Choi YY, Park K, Kim MJ, Kim SH. Neuroregenerative potential of intravenous G-CSF and autologous peripheral blood stem cells in children with cerebral palsy: A randomized, double-blind, cross-over study. *J Transl Med*

- 2017; **15**: 16 [PMID: [28109298](#) DOI: [10.1186/s12967-017-1120-0](#)]
- 43 **Rosenbaum PL**, Walter SD, Hanna SE, Palisano RJ, Russell DJ, Raina P, Wood E, Bartlett DJ, Galuppi BE. Prognosis for gross motor function in cerebral palsy: Creation of motor development curves. *JAMA* 2002; **288**: 1357-1363 [PMID: [12234229](#) DOI: [10.1001/jama.288.11.1357](#)]
- 44 **Luan Z**, Liu W, Qu S, Du K, He S, Wang Z, Yang Y, Wang C, Gong X. Effects of neural progenitor cell transplantation in children with severe cerebral palsy. *Cell Transplant* 2012; **21** Suppl 1: S91-S98 [PMID: [22507684](#) DOI: [10.3727/096368912X633806](#)]
- 45 **Chen G**, Wang Y, Xu Z, Fang F, Xu R, Wang Y, Hu X, Fan L, Liu H. Neural stem cell-like cells derived from autologous bone mesenchymal stem cells for the treatment of patients with cerebral palsy. *J Transl Med* 2013; **11**: 21 [PMID: [23351389](#) DOI: [10.1186/1479-5876-11-21](#)]
- 46 **Huang L**, Zhang C, Gu J, Wu W, Shen Z, Zhou X, Lu H. A Randomized, Placebo-Controlled Trial of Human Umbilical Cord Blood Mesenchymal Stem Cell Infusion for Children With Cerebral Palsy. *Cell Transplant* 2018; **27**: 325-334 [PMID: [29637820](#) DOI: [10.1177/0963689717729379](#)]
- 47 **Kang M**, Min K, Jang J, Kim SC, Kang MS, Jang SJ, Lee JY, Kim SH, Kim MK, An SA, Kim M. Involvement of Immune Responses in the Efficacy of Cord Blood Cell Therapy for Cerebral Palsy. *Stem Cells Dev* 2015; **24**: 2259-2268 [PMID: [25977995](#) DOI: [10.1089/scd.2015.0074](#)]
- 48 **Amati E**, Sella S, Perbellini O, Alghisi A, Bernardi M, Chiericato K, Lievore C, Peserico D, Rigno M, Zilio A, Ruggeri M, Rodeghiero F, Astori G. Generation of mesenchymal stromal cells from cord blood: Evaluation of in vitro quality parameters prior to clinical use. *Stem Cell Res Ther* 2017; **8**: 14 [PMID: [28115021](#) DOI: [10.1186/s13287-016-0465-2](#)]
- 49 **Oppliger B**, Joerger-Messerli M, Mueller M, Reinhart U, Schneider P, Surbek DV, Schoeberlein A. Intranasal Delivery of Umbilical Cord-Derived Mesenchymal Stem Cells Preserves Myelination in Perinatal Brain Damage. *Stem Cells Dev* 2016; **25**: 1234-1242 [PMID: [27392671](#) DOI: [10.1089/scd.2016.0027](#)]
- 50 **Mueller M**, Oppliger B, Joerger-Messerli M, Reinhart U, Barnea E, Paidas M, Kramer BW, Surbek DV, Schoeberlein A. Wharton's Jelly Mesenchymal Stem Cells Protect the Immature Brain in Rats and Modulate Cell Fate. *Stem Cells Dev* 2017; **26**: 239-248 [PMID: [27842457](#) DOI: [10.1089/scd.2016.0108](#)]
- 51 **Hanna SE**, Bartlett DJ, Rivard LM, Russell DJ. Reference curves for the Gross Motor Function Measure: Percentiles for clinical description and tracking over time among children with cerebral palsy. *Phys Ther* 2008; **88**: 596-607 [PMID: [18339799](#) DOI: [10.2522/ptj.20070314](#)]
- 52 **Fleiss B**, Gressens P. Tertiary mechanisms of brain damage: A new hope for treatment of cerebral palsy? *Lancet Neurol* 2012; **11**: 556-566 [PMID: [22608669](#) DOI: [10.1016/S1474-4422\(12\)70058-3](#)]
- 53 **Schoeberlein A**, Mueller M, Reinhart U, Sager R, Messerli M, Surbek DV. Homing of placenta-derived mesenchymal stem cells after perinatal intracerebral transplantation in a rat model. *Am J Obstet Gynecol* 2011; **205**: 277.e1-277.e6 [PMID: [22071064](#) DOI: [10.1016/j.ajog.2011.06.044](#)]
- 54 **Rizvanov AA**, Kiyasov AP, Gaziziov IM, Yilmaz TS, Kaligin MS, Andreeva DI, Shafigullina AK, Guseva DS, Kiselev SL, Matin K, Palotás A, Islamov RR. Human umbilical cord blood cells transfected with VEGF and L(1)CAM do not differentiate into neurons but transform into vascular endothelial cells and secrete neuro-trophic factors to support neuro-genesis-a novel approach in stem cell therapy. *Neurochem Int* 2008; **53**: 389-394 [PMID: [18948156](#) DOI: [10.1016/j.neuint.2008.09.011](#)]
- 55 **Sarnowska A**, Braun H, Sauerzweig S, Reymann KG. The neuroprotective effect of bone marrow stem cells is not dependent on direct cell contact with hypoxic injured tissue. *Exp Neurol* 2009; **215**: 317-327 [PMID: [19063882](#) DOI: [10.1016/j.expneurol.2008.10.023](#)]



Published By Baishideng Publishing Group Inc
7041 Koll Center Parkway, Suite 160, Pleasanton, CA 94566, USA
Telephone: +1-925-2238242
E-mail: bpgoffice@wjgnet.com
Help Desk: <https://www.f6publishing.com/helpdesk>
<https://www.wjgnet.com>

